

ERRATUM

In the manuscript “Mental health of public health workers in Bento Gonçalves, Rio Grande do Sul, Brazil”, DOI: 10.5327/Z1679443520180196, published in the *Revista Brasileira de Medicina do Trabalho*, 2018;16(2):145-57.

Where it reads:

Table 4. Multivariate analysis: factors associated with common mental disorders (CMD) and suicidal ideation, Bento Gonçalves, 2017 (n=567).

Variables	CMD		Suicidal ideation	
	Crude PR (CI)	Adjusted PR (CI)	Crude PR (CI)	Adjusted PR (CI)
Sex	p=0.09	p=0.17	p=0.10	p=0.28
Male	1	1	1	1
Female	1.60 (0.93-2.75)	1.45 (0.85-2.49)	1.88 (0.88-3.99)	1.52 (0.72-3.24)
Age range (years old)	p=0.05	p=0.03	p=0.82	----
Up to 25	1	1	1	
26 to 35	0.55 (0.35-0.88)	0.53 (0.27-0.83)	0.81 (0.41-1.62)	
35 to 45	0.57 (0.34-0.95)	0.48 (0.27-0.86)	0.85 (0.40-1.77)	
46 and over	0.58 (0.35-0.96)	0.47 (0.37-0.94)	1.04 (0.51-2.12)	
Formal schooling	p=0.70	-----	p=0.004*	p=0.002*
11 years (secondary school)	1		1	1
12 to 15 years	0.93 (0.61-1.41)		0.49 (0.28-0.83)	0.50 (0.28-0.89)
16 years or more	1.12 (0.74-1.70)		0.43 (0.23-0.81)	0.44 (0.23-0.83)
Marital status	p=0.72	----	p=0.05	p=0.07
Married/with partner	1		1	1
Single	1.06 (0.71-1.56)		0.82 (0.47-1.42)	0.95 (0.53-1.70)
Separated/widowed	1.26 (0.72-2.22)		1.87 (1.02-3.40)	1.89 (1.06-3.38)
Permanent civil servant	p=0.08	p=0.02	p=1.00	-----
No	1	1	1	
Yes	1.38 (0.97-1.97)	1.60 (1.08-2.39)	1.00 (0.64-1.56)	
Smoking	p=0.42	-----	p=0.03	p=0.11
Never	1		1	1
Ex-smoker	1.38 (0.83-2.30)		1.98 (1.08-3.61)	1.69 (0.89-3.21)
Smoker	1.18 (0.73-1.93)		1.71 (0.98-2.98)	1.63 (0.85-2.81)
Length in job	p=0.005	p=0.007	p=0.59	-----
Up to 11 months	1	1	1	
1 to <4 years	2.37 (1.39-4.04)	2.25 (1.29-3.92)	1.56 (0.80-3.04)	
4 to <9 years	1.60 (0.86-2.97)	1.36 (0.69-2.70)	1.52 (0.74-3.11)	
9 years or more	1.34 (0.73-2.47)	1.16 (0.54-2.51)	1.32 (0.66-2.67)	

Continue...

DOI: 10.5327/Z1679443520180196erratum

Table 4. Continuation.

Variables	CMD		Suicidal ideation	
	Crude PR (CI)	Adjusted PR (CI)	Crude PR (CI)	Adjusted PR (CI)
Professional category	p<0.001	p<0.001	p<0.001	p=0.01
Nursing technicians/assistants	1	1	1	1
Dental, radiology, laboratory technicians	1.39 (0.48-4.06)	1.03 (0.24-4.45)	1.03 (0.37-2.90)	1.01 (0.39-2.63)
Physicians	1.39 (0.63-3.07)	2.48 (1.03-5.95)	0.50 (0.19-1.33)	0.48 (0.15-1.57)
Nurses	1.77 (0.82-3.83)	1.56 (0.70-3.49)	0.46 (0.16-1.32)	0.40 (0.13-1.26)
Other high level professionals	1.68 (0.76-3.69)	1.72 (0.82-3.62)	--- ^a	--- ^a
Administrative employees	1.68 (0.83-3.37)	2.05 (1.00-4.20)	0.73 (0.35-1.50)	0.70 (0.32-1.52)
Inspectors/surveillance	3.16 (1.19-8.39)	2.43 (0.75-7.85)	2.50 (0.98-6.38)	2.26 (0.74-6.89)
Cleaning	3.03 (1.35-6.79)	3.84 (1.63-9.04)	2.11 (1.01-4.41)	1.70 (0.80-3.64)
CHA	3.96 (2.03-7.70)	7.43 (3.28-16.84)	1.94 (1.00-3.78)	1.88 (0.95-3.75)
Conflict with supervisors	p<0.001	p<0.001	p=0.004	P<0.001
No	1	1	1	1
Yes	2.65 (1.85-3.80)	3.13 (2.15-4.56)	2.10 (1.26-3.51)	2.40 (1.47-3.93)
Conflict with coworkers	p<0.001	p<0.001	p=0.001	p<0.001
No	1	1	1	1
Yes	3.45 (2.45-4.84)	3.05 (2.11-4.40)	2.22 (1.36-3.62)	2.56 (1.57-4.20)
Conflict with service users	p<0.001	p=0.001	p=0.02	p=0.002
No	1	1	1	1
Yes	2.12 (1.47-3.05)	1.85 (1.28-2.68)	1.84 (1.11-3.06)	2.15 (1.34-3.47)

PR: prevalence ratio; CI: 95% confidence interval; ACS: community health agents; *linear trend estimation; only associations with p=0.20 were subjected to regression analysis; CMD adjusted for sex, age range, educational level, employment relationship, length of work in current job, professional category; suicidal ideation adjusted for sex, educational level, marital status, smoking and professional category; ^aexcluded from analysis due to low number of participants.

It should read:

Table 4. Multivariate analysis: factors associated with common mental disorders (CMD) and suicidal ideation, Bento Gonçalves, 2017 (n=567).

Variables	CMD		Suicidal ideation	
	Crude PR (CI)	Adjusted PR (CI)	Crude PR (CI)	Adjusted PR (CI)
Sex	p=0.09	p=0.17	p=0.10	p=0.28
Male	1	1	1	1
Female	1.60 (0.93-2.75)	1.45 (0.85-2.49)	1.88 (0.88-3.99)	1.52 (0.72-3.24)
Age range (years old)	p=0.05	p=0.03	p=0.82	----
Up to 25	1	1	1	
26 to 35	0.55 (0.35-0.88)	0.53 (0.27-0.83)	0.81 (0.41-1.62)	
35 to 45	0.57 (0.34-0.95)	0.48 (0.27-0.86)	0.85 (0.40-1.77)	
46 and over	0.58 (0.35-0.96)	0.47 (0.37-0.94)	1.04 (0.51-2.12)	

Continue...

Table 4. Continuation.

Variables	CMD		Suicidal ideation	
	Crude PR (CI)	Adjusted PR (CI)	Crude PR (CI)	Adjusted PR (CI)
Formal schooling	p=0.70	-----	p=0.004*	p=0.002*
Up to 11 years (secondary school)	1		1	1
12 to 15 years	0.93 (0.61-1.41)		0.49 (0.28-0.83)	0.50 (0.28-0.89)
16 years or more	1.12 (0.74-1.70)		0.43 (0.23-0.81)	0.44 (0.23-0.83)
Marital status	p=0.72	---	p=0.05	p=0.07
Married/with partner	1		1	1
Single	1.06 (0.71-1.56)		0.82 (0.47-1.42)	0.95 (0.53-1.70)
Separated/widowed	1.26 (0.72-2.22)		1.87 (1.02-3.40)	1.89 (1.06-3.38)
Permanent civil servant	p=0.08	p=0.02	p=1.00	-----
No	1	1	1	
Yes	1.38 (0.97-1.97)	1.60 (1.08-2.39)	1.00 (0.64-1.56)	
Smoking	p=0.42	-----	p=0.03	p=0.11
Never	1		1	1
Ex-smoker	1.38 (0.83-2.30)		1.98 (1.08-3.61)	1.69 (0.89-3.21)
Smoker	1.18 (0.73-1.93)		1.71 (0.98-2.98)	1.63 (0.85-2.81)
Length in job	p=0.005	p=0.007	p=0.59	-----
Up to 11 months	1	1	1	
1 to <4 years	2.37 (1.39-4.04)	2.25 (1.29-3.92)	1.56 (0.80-3.04)	
4 to <9 years	1.60 (0.86-2.97)	1.36 (0.69-2.70)	1.52 (0.74-3.11)	
9 years or more	1.34 (0.73-2.47)	1.16 (0.54-2.51)	1.32 (0.66-2.67)	
Professional category	p<0.001	p<0.001	p<0.001	p=0.01
Nursing technicians/assistants	1	1	1	1
Dental, radiology, laboratory technicians	1.39 (0.48-4.06)	1.03 (0.24-4.45)	1.03 (0.37-2.90)	1.01 (0.39-2.63)
Physicians	1.39 (0.63-3.07)	2.48 (1.03-5.95)	0.50 (0.19-1.33)	0.48 (0.15-1.57)
Nurses	1.77 (0.82-3.83)	1.56 (0.70-3.49)	0.46 (0.16-1.32)	0.40 (0.13-1.26)
Other high level professionals	1.68 (0.76-3.69)	1.72 (0.82-3.62)	--- ^a	--- ^a
Administrative employees	1.68 (0.83-3.37)	2.05 (1.00-4.20)	0.73 (0.35-1.50)	0.70 (0.32-1.52)
Inspectors/surveillance	3.16 (1.19-8.39)	2.43 (0.75-7.85)	2.50 (0.98-6.38)	2.26 (0.74-6.89)
Cleaning	3.03 (1.35-6.79)	3.84 (1.63-9.04)	2.11 (1.01-4.41)	1.70 (0.80-3.64)
CHA	3.96 (2.03-7.70)	7.43 (3.28-16.84)	1.94 (1.00-3.78)	1.88 (0.95-3.75)
Conflict with supervisors	p<0.001	p<0.001	p=0.004	P<0.001
No	1	1	1	1
Yes	2.65 (1.85-3.80)	3.13 (2.15-4.56)	2.10 (1.26-3.51)	2.40 (1.47-3.93)
Conflict with coworkers	p<0.001	p<0.001	p=0.001	p<0.001
No	1	1	1	1
Yes	3.45 (2.45-4.84)	3.05 (2.11-4.40)	2.22 (1.36-3.62)	2.56 (1.57-4.20)
Conflict with service users	p<0.001	p=0.001	p=0.02	p=0.002
No	1	1	1	1
Yes	2.12 (1.47-3.05)	1.85 (1.28-2.68)	1.84 (1.11-3.06)	2.15 (1.34-3.47)

PR: prevalence ratio; CI: 95% confidence interval; ACS: community health agents; *linear trend estimation; only associations with p=0.20 were subjected to regression analysis; CMD adjusted for sex, age range, educational level, employment relationship, length of work in current job, professional category; suicidal ideation adjusted for sex, educational level, marital status, smoking and professional category; ^aexcluded from analysis due to low number of participants.